

Naloxone Pre/Post Test

- 1) **What can increase your risk of having an overdose? (choose 1 answer)**
 - a) mixing drugs
 - b) using alone
 - c) using after a period of non-use
 - d) all of the above

- 2) **Which of the following are three signs of an opioid overdose?**
 - a) unable to wake the person up
 - b) not breathing at all or breathing very slow
 - c) turning blue/purple around lips and finger tips
 - d) Increase energy, wanting to exercise

- 3) **Is it essential that you call 911 for all overdoses?**
 - a) Yes
 - b) No

- 4) **Does Naloxone work for a cocaine overdose?**
 - a) Yes
 - b) No

- 5) **Why must you stay and support the person that overdosed? (choose 1 answer)**
 - a) naloxone may wear off and overdose may return.
 - b) may need to give a 2nd dose of naloxone
 - c) provide important information to EMS
 - d) All of the above

- 6) **How many cc's of naloxone do you administer for each dose?**
 - a) 5cc
 - b) 1cc
 - c) 10cc
 - d) .5cc

- 7) **How long does it take for naloxone to start working once given IM (intra muscular)?**
 - a) 10 min
 - b) 1-5 minutes
 - c) 20 minutes

- 8) **How many doses of naloxone do you administer for an opioid overdose?**
 - a) 10 doses
 - b) Start with 1 dose, if no response in 3 – 5 mins give 2nd dose, ensure 911 has been called
 - c) 1 dose